

THE EVOLUTION OF TYPE 1 AGN AT 15 μm FROM THE ELAIS

F. LA FRANCA¹, I. MATUTE¹, C. GRUPPIONI², D. ALEXANDER³, + *ELAIS CONSORTIUM*

¹*Dipartimento di Fisica, Università Roma Tre, Via della Vasca Navale 84, 00146, Roma, Italy*

²*Osservatorio di Padova, Vicolo dell'Osservatorio 35, 35122, Padova, Italy*

³*SISSA, Via Beirut 4, 00000, Trieste, Italy*

ABSTRACT.

We present the first preliminary estimate of the evolution of type 1 AGNs at 15 μm . A new sample of sources selected from the ELAIS-S1 survey has been used together with the RMS sample. The ELAIS-S1 survey covers an *effective* area of 2.2 deg² (total: 4 deg²) down to 1.2 mJy at 15 μm . Spectroscopic identifications have been carried out in the range 17.2 < R < 20.0 at the 3.6m/ESO, NTT/ESO and 2dF/AAT. The luminosity function is steeper at bright luminosities in comparison to the previous estimates of Rush, Malkan & Spinoglio (1993). The evolution is compatible with a Pure Luminosity Evolution model with $L(z) \propto L(0)(1+z)^{3.4}$.

1. The European Large Area ISO Survey (ELAIS)

ELAIS is a collaboration involving 25 European Institutes, led from Imperial College. ELAIS is one of the largest single Open Time project conducted by ISO, mapping an area of 12 squares degrees at 15 μm with ISOCAM and at 90 μm with ISO-PHOT. Secondary surveys in other ISO bands were undertaken by the ELAIS team within the fields of the primary survey, with 6 degrees being covered at 6.7 μm and 1 degree at 175 μm (Oliver et al, 2000). Four main fields were chosen (N1, N2, N3 in the north hemisphere and S1 in the south) at high Ecliptic latitudes ($|\beta| > 40^\circ$).

The work here presented is the result of the follow-up campaign carried out in the field S1 (J2000, $\alpha : 00^h 34^m 44^s$, $\delta : -43^\circ 34' 44''$). An initial catalog was produced by the Imperial College using their data reduction technique ("Preliminary Analysis"). Optical identifications were possible thanks to an extensive R-band CCD survey, performed with the ESO/Danish 1.5m telescope. The spectroscopic follow-up program resulted in ~ 130 redshift identifications with the 2dF/AAT, 3.6/ESO and NTT/ESO. The sample is flux limited at 15 μm down to 1.2 mJy, and in the R-band to magnitudes between 17.2 and 20.0.

2. The luminosity function at 15 μm

In order to compute the luminosity function (LF) of type 1 AGNs (AGN1) at 15 μm , we added to our sample the AGN1 from the catalog of Rush, Malkan & Spinoglio (1993,

RMS) as representative of the local population. The RMS consists of a sample of galaxies selected a $12\ \mu\text{m}$ from the IRAS PSCv2 complete down to 0.3 Jy.

We characterize our sources in the Infrared with a single SED. Ground-based observations (Roche et al. 1991; Moorwood 1986) and the latest ISO observations on Type 1 AGNs find that the Mid-Infrared (MIR) broadband emission features (from PAH molecules) are generally absent in this kind of sources (Lutz et al. 1997a; Genzel et al. 1998). For this reason the mean SED for radio-quiet Quasars from Elvis et al. (1994) was chosen. The standard K-correction (in magnitudes) was computed following Lang (1980). In order to compute the $15\ \mu\text{m}$ luminosities and K -corrections, we convolved the SED from Elvis et al. (1994) with the IRAS $12\ \mu\text{m}$ and ISOCAM $15\ \mu\text{m}$ bandpasses.

We assumed a simple double power-law luminosity function (LF) and pure luminosity evolution (PLE) of the type:

$$\begin{aligned}\Phi(L) &= CL_*^{(\alpha-\beta)} L^{-\alpha} & \text{for } L \leq L_*, \\ \Phi(L) &= CL^{-\beta} & \text{for } L > L_*\end{aligned}$$

where α and β are the slopes of the faint and bright end of the LF, L_* the luminosity of the 'knee' of the LF and C the normalization factor. The evolution in luminosity is given by the relation:

$$L_*(z) = L_*(0) (1+z)^\kappa.$$

A parametric, unbinned maximum likelihood method was used to fit the evolution and luminosity function parameters simultaneously (Marshall et al. 1983) at $15\ \mu\text{m}$. Since ELAIS identifications were not only flux limited at $15\ \mu\text{m}$, but also in their R-band magnitude ($17.2 < R < 20.0$), a factor Θ was introduced in the function 'S' to be minimized

$$S = -2 \sum_{i=1}^N \ln[\Phi(z_i, L_i)] + \iint \Phi(z, L) \Omega(z, L) \Theta(z, L) \frac{dV}{dz} dz dL$$

and only applied to the ELAIS sample. This factor represents the conditional probability of a source with a given $15\ \mu\text{m}$ luminosity to have R magnitude in the range (17.2,20.0):

$$\Theta(17.2 < R < 20.0 \mid L_{15}),$$

computed using the local (RMS) L_R/L_{15} distribution, supposed constant with z .

3. Results and comments

For the first time, the evolution of the Type 1 AGNs luminosity function at $15\ \mu\text{m}$ is derived from a list of sources selected in the Infrared (Fig.1). The LF evolves according to a PLE model with $\kappa \simeq 3.4$, an evolution similar to the one derived in the optical domain (e.g. La Franca & Cristiani, 1997).

The slope of the previous derivation of the LF from RMS has a flatter slope in the bright part ($\beta = 2.1$) in comparison with our determination ($\beta = 2.5$). This overestimate of the density of bright AGN1 by RMS is due to the effects of evolution inside the redshift bins used with the $1/V_a$ method.




Fig. 1. Best-fitting PLE model for 2 redshifts intervals (*solid lines*). The points (*rhombuses, squares*) correspond to the space-densities of the observed sources, corrected for evolution within the redshift intervals. As a comparison, the RMS fit of the luminosity function (*dashed line*) and our LF at $z=0$ (*dots*) are also plotted.

References

- Elvis, M., Wilkes, B.J., McDowell, J.C., et al. 1994, *Astrophys. J. Suppl.* **95**, 1.
 Genzel, R. et al. 1998, *Astrophys. J.* **498**, 579.
 La Franca, F., Cristiani, S. 1997, *Astron. J.* **113**, 1517.
 Lang, K.R. 1980, *Astrophysical Formulae* (Berlin:Springer).
 Lutz, D. et al. 1997, *Astron. Astrophys. Suppl. Ser.* **248**, 217.
 Marshall, H.L., Avni, Y., Tananbaum, H., Zamorani, G. 1983, *Astrophys. J.* **269**, 35.
 Moorwood, A.F. 1986, *Astron. Astrophys. Lett.* **137**, L5.
 Oliver, S. et al, *Mon. Not. R. Astr. Soc.* ., in press, astro-ph/0003263.
 Roche, P.F., Aitken, D.K., Smith, C.H., & Martin J.W. 1991, *Mon. Not. R. Astr. Soc.* **248**, 606.
 Rush, B., Malkan, M.A. & Spinoglio, L. 1993, *Astrophys. J. Suppl.* **89**, 1.
 Xu, C., Hacking, P.B., Fang, F., Shupe, D.L., Lonsdale, C., Lu, N.Y., Helou, G. 1998, *Astrophys. J.* **508**, 576.